REPORT DOCUMENTATION PAGE	Form Approved OMB No. 0704-0188
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing maintaining the data peeded, and completing and reviewing this collection of information.	instructions, searching existing data sources, gathering and
Highway, Suite 1204 Arlington, VA 22202-4302 Respondents should be away that the adjustness Services, Directorate for information of the services of the servi	ation Operations and Reports (0704-0188), 1215 Jefferson Davis
collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO T 1. REPORT DATE (DD-MM-YYYY) 2. REPORT TYPE	HE ABOVE ADDRESS.
Technical Papers	3. DATES COVERED (From - To)
4. TITLE AND SUBTITLE	5a. CONTRACT NUMBER
	5b. GRANT NUMBER
·	5c. PROGRAM ELEMENT NUMBER
6. AUTHOR(S)	5d. PROJECT NUMBER
	2303
	5e. TASK NUMBER
	<u>m 208</u>
	5f. WORK UNIT NUMBER
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)	8. PERFORMING ORGANIZATION
Air Force Research Laboratory (AFMC)	REPORT
AFRL/PRS	
5 Pollux Drive	
Edwards AFB CA 93524-7048	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)	10.000.000
AND ADDITION (C)	10. SPONSOR/MONITOR'S ACRONYM(S)
Air Force Research Laboratory (AFMC)	i
AFRL/PRS	11 CDONCODITIONITODIO
5 Pollux Drive	11. SPONSOR/MONITOR'S NUMBER(S)
Edwards AFB CA 93524-7048	
12. DISTRIBUTION / AVAILABILITY STATEMENT	· ·
	*
Approved for public release; distribution unlimited.	
13. SUPPLEMENTARY NOTES	
14. ABSTRACT	
	•
5. SUBJECT TERMS	
129	020
	IUMBER 19a. NAME OF RESPONSIBLE
OF ABSTRACT OF P	PAGES PERSON
REPORT b. ABSTRACT c. THIS PAGE	Leilani Richardson 19b. TELEPHONE NUMBER
Unclassified Unclassified Unclassified (A)	(include area code)
Unclassified Unclassified Unclassified	(661) 275-5015
	Standard Form 298 (Rev. 8-98) Prescribed by ANSI Std. 239.18

MEMORANDUM FOR PRS

FROM: PROI (TI) (STINFO)

28 May 1999

SUBJECT: Authorization for Release of Technical Information, Control Number: AFRL-PR-ED-TP-FY99-0117 Yoo and Boatz, "Theoretical Study of Decomposition Mechanism of High Energy Density Materials"

Poster Session HEDM Conference

(Statement A)

20021122 020

Theoretical Study of Decomposition Mechanism of High Energy Density Materials

Hi Young Yoo and Jerry Boatz*

*Air Force Research Laboratory, Edwards AFB, CA 93524-7680 National Research Council, Washington, DC 20418

AFOSR HEDM CONTRACTOR'S MEETING 6/8-6/10, 1999
Cocoa Beach, FL

DISTRIBUTION STATEMENT A: Approved for Public Release -Distribution Unlimited

Abstract

Decomposition mechanisms of [NH₂Me₂]⁺[NO₃], and [NH₂(NH₂)₂]⁺[NO₃] were processes have been explored: (1) proton transfer and (2) Me⁺/NH₂⁺ transfer 31G* level and single point energies at the MP2/6-31G*//RHF/6-31G* level have been calculated. Comparison of similarities and differences between these two systems will be presented here. Future research will involve characterization of the potential energy surface for [NH₂NMe₂NH₂]⁺[NO₃] and design of potential catalysts One of the HEDM research efforts has focused on the development of high investigated. The potential energy surfaces of two gas-phase decomposition reactions. Transition states for both pathways have been located at the RHF/6proposed area of research involves determination of decomposition mechanisms and stabilities of monopropellants by using ab initio quantum mechanical calculations. performance monopropellants in replacement for hydrazine due to its toxicity. A that will stabilize transition states.

$$[NH_2Me_2]^+[NO_3]^- \rightarrow NHMe_2 + HNO_3$$
 (1a)

$$[NH_2Me_2]^+[NO_3]^- \rightarrow NH_2Me + MeNO_3$$
 (2a)

$$[NH_2(NH_2)_2]^+[NO_3]^- \to NH(NH_2)_2 + HNO_3$$
 (1b)

$$[NH_2(NH_2)_2]^+[NO_3]^- \to NH_2NH_2 + NH_2NO_3$$
 (2b)

Background

Hydrazine is the state of the art monopropellant currently. However, it has several disadvantages including toxicity, volatility, and handling. Researchers are continually looking for new monopropellant candidates which include energetic materials such as substituted ammonium salts. Our research effort has concentrated on the decomposition routes of some HEDM materials. Ab initio quantum mechanical calculations have been carried out by our group to explore possible decomposition mechanisms.

The model compounds under study are $[NH_2Me_2]^+[X]^-$, where $X = NO_3$ and CI_3 , and $[N(NH_2)_2Me_2]^+[NO3]^-$.

Objectives:

- To map out the potential energy surface of these model compounds
- To determine the energetics of decomposing species
- To design a catalyst that stabilizes the transition state

Gaseous products detected from experimental thermal decomposition studies of $N(NH_2)_2(Me_2)NO_3$ are N_2 , NH_3 , H_2O and CH_4 .

Possible reaction pathways leading to the final products are the following: R, R' = H, CH₃; $X = NO_3$

(1) $[N(NH_2)_2RR']^+[X]^- \rightarrow [N(NH_2)_2R'][RX]$ (1b) $[N(NH_2)_2RR']^+[X]^- \rightarrow [N(NH_2)RR'][NH_2X]$

(2a) $N(NH_2)_2R' \rightarrow N(NH_2)R' + NH_2$ (2b) $N(NH_2)RR' \rightarrow NRR' + NH_2$ (3) $[N(NH_2)_2RR']^+ \rightarrow RH + [NH_3R']^+ + N_2$

(4a) $NH_2X \rightarrow NH_2 + X$

4b) $NH_2 + NH_2X \rightarrow NH_3 + NHX$

(4c) $X + NH_2X \rightarrow HX + NHX$

(4d) $2NHX \rightarrow X-NH-NH-X$

(4e) X-NH-NH-X \rightarrow 2HX + N₂

Rubtsov, Y. I.; Andrienko, L. P.; Titova, K. V.; Loginova, E. N. Izvestiya Akademii Nauk SSSR, Seriya Khimicheskaya 1982, 9, 1953.

Decomposition Mechanisms of [NH₂(NH₂)₂]⁺[X]⁻, $X = CI \text{ or } NO_3$


Proton Transfer


$$[NH_2(NH_2)_2]^+[X]^- \rightarrow NH(NH_2)_2 + HX$$

NH2 Cation Transfer

$$[NH_2(NH_2)_2]^+[X]^- \rightarrow NH_2NH_2 + NH_2X$$

A Proton Transfer Transition Structure of [NH₂(NH₂)₂]⁺[NO₃] At RHF/6-31G*(6d)





Intermediate Complex


Energy 0.0 (kcal/mol) (0.0)


Transition Structure

7

(-1.2)

A NH2⁺ Transfer Transition Structure of [NH2(NH2)2]⁺[NO3] At RHF/6-31G*(6d)





Intermediate Complex

0.0 (0.0)


(kcal/mol)

Energy


Transition Structure


8.69

(55.9)



A Proton Transfer Transition Structure of [NH2Me2] [NO3] At RHF/6-31G*(6d)





Intermediate Complex


(kcal/mol) Energy


0.0 (0.0)

Transition Structure

2.2 (-1.3)

A Me⁺ Transfer Transition Structure of [NH₂Me₂]⁺[NO₃]⁻ At RHF/6-31G*(6d)






Intermediate Complex

Energy (kcal/mol)

0.0 (0.0)

Transition Structure

45.8 (47.7)



Conclusion

Based on the ab initio calculation studies of [NH₂(NH₂)₂]⁺[NO₃ or Cl] and $[\mathrm{NH}_2(\mathrm{Me})_2]^{+}[\mathrm{NO}_3 \text{ or CI}]^{-}$

- The transition structure of proton transfer is earlier than that of methyl(or NH_2) cation transfer and also lower in barrier.
- Variations on anions or cations did not affect on the energy barrier or the transition structure.
- calculations imply that methyl cation transfer of [NMe₂(NH₂)₂]⁺[NO₃] will Since variations on cations did not affect on the energy barrier, our be favorable than NH₂⁺.

Future Work

- The decomposition mechanism of $[NMe_2(NH_2)_2]^+[NO_3]^-$ as well as other ammonium salt monopropellant candidates will be investigated theoretically.
- potential energy surface of [NMe₂(NH₂)₂]⁺[NO₃] and it will be compared Different catalysts will be included in the calculations for determining the with the uncatalyzed potential energy surface.